Science 10 Mutations and DNA




GENETICS

Part I- The Nucleus: Control Centre of the Cell


 Every cell in your body has a specific <u>JOB</u>- but how do they become specialized?
 E.g. hair cells vs. skin cells vs. retina in the eye

The **NUCLEUS** in the cell contains the master set of instructions that tells the cell:


- what it will **BECOME**
- how it will function
- when it will <u>REPRODUCE</u> and <u>GROW</u>
- when it will die



• But how does the nucleus do this? How does it send messages to the rest of the cell?




The nucleus contains <u>DNA</u>, which carries the master set of <u>INSTRUCTIONS</u> for cell function.



 DNA (<u>DEOXYRIBONUCLEIC ACID</u>) is a double stranded helix that looks like a twisted ladder

- the sides of DNA are made of <u>SUGAR</u> and <u>PHOSPHATES</u>
- the steps of DNA are made up of 4 BASES:

AdenineCytosineGuanineThymine



- Where are the bases located?
- Name the 4 bases
- Where are the sugars and phosphates located?
- What is this shape called?




Figure 4.5 The structure of the DNA molecule

Pairing Rules

- adenine (A) always pairs with <u>THYMINE</u> (T)
 → Apple Tree
- cytosine (C) always pairs with <u>GUANINE</u> (G)
 → Car Garage





2. DNA has many functions:

A. DNA contains **INSTRUCTIONS FOR ALL CELL FUNCTIONS** and, therefore, DNA indirectly controls all of the functioning of all living things.

B. DNA **DETEMINES THE HEREDITARY TRAITS** of an individual


- C. DNA <u>EVOLVES</u> (changes through mutations and recombination). This allows for new characteristics & abilities to appear which may help an individual to survive & reproduce.
- D. Self replication: DNA has the ability to **MAKE COPIES OF ITSELF.**

3. The arrangement of bases in DNA directs all cell activity

- the bases are like letters that carry a message (CODE)
- the code gives **INSTRUCTIONS** for a specific task.




4. DNA is stored in the form of chromatin




- chromatin is made up of DNA and **PROTIENS** called histones (very dense)
- when a cell is growing, parts of the <u>CHROMATIN</u> unwind so that the targeted section of DNA can be read to make messages that control the rest of the cell.
- when cells reproduce, the entire chromatin coils up and makes an X-shaped structure called a <u>CHROMOSOME</u>


Some terms you should know....



5. Every organism has a characteristic number of chromosomes


- chromosomes are always in **PAIRS** in the nucleus
- humans have 46 chromosomes (=<u>23</u> pairs)
- in males, the 23rd pair of chromosomes are "XY"
- in females, the 23rd pair of chromosomes are "XX"
- cows have 60 chromosomes; corn has 20 chromosomes






6. Genes are found on chromosomes

- A <u>GENE</u> is a small segments of DNA found at specific places on a chromosome that code for a protein
- genes can vary in length from 100s to 1000s of <u>BASES</u>
- the arrangement of bases will decide what kind of protein is produced



- each chromosome carries 1000s of <u>GENES</u>
- your body uses 90 000 to 100 000 different
 PROTEINS

https://www.youtube.com/watch?v=7Hk9jct2ozY




brown eyes.

Part I- The Role of Proteins

Why are proteins so important anyways?

- Humans share most of the same protein families with <u>WORMS</u>, flies, and plants
- Hair grows by forming new cells at the base of the root. As they
 move upward through the skin they are cut off from their
 nutrient supply and start to form a hard protein called <u>KERATIN.</u>
 As this occurs, the hair cells die. The dead cells and keratin form
 the shaft of the hair.

- Fingernails grow about **THREE TO FOUR** times as quickly as toenails
- Each hair grows about <u>1/4 INCH</u>/month and grows for up to <u>6</u> years.
- The most expressive muscles are the facial muscles. We need <u>17</u> muscles to smile and <u>43</u> muscles to frown.
- The most numerous are the skeletal muscles. When we walk for instance, we use <u>200</u> muscles.



Proteins Have 2 Main Functions

1. Structural: proteins help make up all structures in living things
 Examples: a) <u>MUSCLE PROTEINS</u>
 b) <u>HAIR, NAILS, BONES</u>

c) **BLOOD VESSELS, LIGAMENTS**

2. Functional: other proteins help us to keep our bodies functioning properly and to digest our food.

Examples: a) <u>ENZYMES TO SPEED UP CHEMICAL REACTIONS</u> b) <u>HAEOMOGLOBIN IN RED BLOOD CELLS</u> c) <u>ANTIBODIES USED IN THE IMMUNE SYSTEM</u> d) <u>TRANSPORT PROTEINS FOUND IN CELL MEMBRANES</u>


- Only certain genes are <u>ACTIVATED</u> in a cell
- Depending which genes are active, different proteins are produced, and this causes cells to have different functions.

E.x. You do not have skin pigment genes being used by your stomach cells

• ONE GENE \rightarrow ONE TYPE OF PROTEIN \rightarrow ONE FUNCTION


Part II- Protein Synthesis

- The making of proteins can be broken down into two steps:
 - 1. TRANSCRIPTION
 - 2. TRANSLATION



<u>Act One:</u> <u>TRANSCRIPTION</u> (*Trans* = across, *cription* = to write)


- The coded message of a gene on DNA has INSTRUCTIONS on how to make particular <u>PROTEINS</u> that our bodies need.
- The instructions from a gene are copied from DNA to <u>MESSANGER RIBONUCLEIC ACID</u> (MRNA) in the nucleus.



- RNA has slightly different bases: A, G, C and **URACIL** instead of thymine
- Once complete, the mRNA moves through the <u>NUCLEAR PORES</u> and into the cytoplasm where the proteins are made.
- The process of making mRNA is called **TRANSCRIPTION**

Watch the General Process


- **Step 1**: <u>UNZIPPING</u> of the DNA from a double strand into two single strands
- **Step 2**: <u>**RIBOSOMAL BASES**</u> pair up with the bases on one of the DNA strands to form mRNA
- Step 3: ENZYMES help form the RNA BACK BONE and checks for ERRORS
- Step 4: The <u>MRNA</u> is released, leaves the nucleus, & the <u>DNA ZIPS BACK</u> <u>UP</u>



Act Two: **TRANSLATION**

- The mRNA moves into the cytoplasm and pairs up with a **<u>RIBOSOME</u>**. It is here that the mRNA will be "read" to make a protein.
- The mRNA code is made up of groups of <u>TRIPLET_CODES</u> known as <u>CODONS</u>. Each codon codes for a specific <u>AMINO ACID</u>. Eg. AGC = Serine Eg. UGC = Cysteine


- It takes <u>3</u> bases to make one codon. Why? Because there are about <u>20</u> amino acids and we only have 3 <u>BASES</u> in the alphabet.
 - With a _SINGLE_ nucleotide, there are only <u>4</u> possible codes (4¹).
 - For _DOUBLE_ nucleotides, there are only <u>16</u> possible codes (4²).
 - However, for _TRIPLET_




mRNA transcribed code


• The amino acid for each codon is given in the tables

			U		c		A	G		
	U		Phenyl- alanine	UCU UCC		UAU UAC	Tyrosine	UGU UGC		
		UUA UUG	Leucine	UCA UCG	Serine	UAA UAG	Stop codon Stop codon	UGA UGG	Stop codon Tryptophan	A G
er	с	CUU CUC	Leucine	CCU	Proline	CAU CAC	Histidine	CGU CGC CGA CGG	Arginine	U C
lette		CUA CUG		CCA CCG		CAA CAG	Glutamine			A G
First	A	AUU AUC	Isoleucine	ACU ACC	Threonine	AAU AAC	Asparagine	AGU AGC	Serine	U C
ш		AUA	Methionine; initiation codon	ACA ACG		AAA AAG	Lysine	AGA AGG	Arginine	A G
	G	GUU GUC	Vallas	GCU GCC	Alesias	GAU GAC	Aspartic acid	GGU GGC	Glycine	U C
		GUA GUG	Valine	GCA GCG	Alanine	GAA GAG	Glutamic	GGA GGG		A G




- During translation, the written code (codons) on mRNA is '<u>TRANSLATED</u>' into a specific amino acid sequence by <u>TRANSFER RIBONUCLUEIC ACID</u> (tRNA) in the cytoplasm.
- A tRNA molecule is a small piece of RNA that have an <u>AMINO ACID</u> attached to it.
- The tRNA also has a special sequence of 3 bases known as an <u>ANTICODON</u>.






- As the correct <u>AMINO ACIDS</u> are brought to the ribosome by the <u>tRNAs</u>, they are joined together to form the <u>PROTEIN CHAIN</u> that the original DNA coded for.
- Eventually, the finished protein chain moves to the **GOLGI BODY**, where it is packaged and released to do its "job".



SUMMARY OF PROTEIN SYNTHESIS:








Figure 4.11 The production of a protein in an animal cell





Worksheet:

DNA AND PROTEIN SYNTHESIS REVIEW

Section 4.2: Mutations

• Real or Fake?



A BAD NIGHT AT THE THEATRE

Question: What if something goes wrong during *translation*?


Answer: IF TRANSLATION IS ALTERED, THE PROTIEN PRODUCT WILL BE DIFFERENT

PROTEIN FUNCITON DEPENDS ON SHAPE! IF THE SHAPE CHANGES, IT'S FUNCTION CHANGES TOO!


- Mutations can <u>CHANGE</u> in the base sequence of DNA
- When the <u>BASES</u> ('letters') change, the <u>INSTRUCTIONS</u> used to make the protein also change.
- Different <u>AMINO ACIDS</u> will be used, resulting in a different protein. This may alter protein shape, and also change or "break" protein function.

There are 2 types of MUTATION:


- 1. <u>CHROMOSOMAL</u> mutations: a mutation of all or part of a chromosome.
- This usually involves <u>LARGE SEGMENTS OF DNA</u>, and therefore, <u>CAN</u> <u>HAVE GREATER IMPACT THAN SMALLER GENE MUTATIONS</u>.
- There are four types of chromosomal mutations that involve part of the chromosome:



Chromosomal Mutations...



- E.g. Fruit fly with legs in place of antennae. This mutation involves moving a controller gene telling the fruit fly where to put the legs
- E.g. Down's syndrome. This is a genetic disorder where there is an extra chromosome #21. The extra chromosome resulted from improper cell division.






Down syndrome karyotype

30 36 36

		,		,	5	,			T			_			
1	2	3 3 20	4	5 00 10 10 10 10 10 10 10 10 10 10 10 10	16	7 7 16	8 17	9			Test Rain Res	Note of the second		flattened nose and face, upward slanting eyes,	
	Norm	al pal	m		Sim			Y		B	ALC C	widely separated first and second toes and increase skin creases	ed		

2. <u>GENE</u> mutations: a mutation that occurs within a gene at some **point** along a chromosome. This mutation is only a change of one or a few 'letters' (bases).

It usually only affects <u>ONE GENE</u>, and therefore, <u>CAN HAVE LESS</u>
 <u>IMPACT THAN LARGER CHROMOSOMAL MUTATIONS</u>.



Recall the sequence of protein synthesis:
 DNA → RNA → protein
 ACTGCT.... UGACGA.... leu- ade-cyt...
 (in nucleus) (to cytoplasm) (at ribosome)



- If the sequence of DNA changes by mutation, then the RNA sequence copied from the DNA will be different, and this will code for a different amino acid, which results in a different protein. This is called a <u>GENE</u> <u>MUTATION</u>
- E.g. the Land of the Spirit Bear is **WHITE** because the gene that codes for **FUR COLOUR** (dark brown) is mutated and no longer works


- there are three main ways that gene mutations can occur:
- DELETION AND ADDITION MOST HARMFUL \rightarrow FRAMESHIFT

Substitution	e.g. replace "C" with "A"	ACTGCA → AATGCA
Deletion	e.g. lose a base all together	ACTGCA → ACGCA
Addition	e.g. add another "A"	ACTGCA → ACTAGCA

I. Types of Gene Mutations


- mutations are often a bad thing because the protein does not work and your body needs it to function. These are known as <u>NEGATIVE</u> <u>MUTATIONS</u> and they <u>DECREASE</u> survival rates.
- e.g. mutated gene \rightarrow **SICKLE-CELL ANEMIA** (misshapen red blood cells that don't carry O₂).
- e.g. more than 1300 different mutations in one gene → <u>CYSTIC FIBROSIS</u> (build up of mucus in the lungs)

Sickle Cell Anemia and Cystic Fibrosis




- sometimes mutations can be a good thing (e.g. the mutated protein works better than the "normal" protein or has a new beneficial function). These are known as <u>POSITIVE</u>
 <u>MUTATIONS</u> and they <u>INCREASE</u> survival rate.
- e.g. mutated gene \rightarrow proteins that make them <u>**RESISTANT**</u> <u>**TO HIV INFECTION**</u>. These people cannot get AIDS.
- e.g. some plants have mutated genes that codes for proteins that make them <u>RESISTANT TO CHEMICAL</u> <u>HERBICIDES</u>.





 sometimes mutations have no effect because they occur in a part of the DNA that is not used for protein synthesis or the mutated protein is not required for survival. These are known as <u>NEUTRAL MUTATIONS</u> and have they have <u>NO EFFECT</u> on the organism.

e.g. mutation in the gene that codes for fur colour (Spirit Bear). Although white, the survival rate of the organism is not changed.



II. Causes of Mutations

- Mutations can occur naturally during DNA replication and cell division. The human body has systems in place that will often fix errors in DNA replication, or destroy cells that are working abnormally due to mutations.
- There are also factors in the environment that may increase chances for mutations to occur.

• <u>MUTAGENS</u>= substances or factors (e.g. virus, cigarette smoke, X-rays) that can cause mutations in DNA

